

Human remains from Ali Kosh, Iran, 2017

Arkadiusz Sołtysiak^{*1}, Hojjat Darabi²

¹ Department of Bioarchaeology,
Institute of Archaeology, University of Warsaw,
Krakowskie Przedmieście 26/28, 00-927 Warsaw, Poland
email: a.soltysiak@uw.edu.pl (corresponding author)

² Department of Archaeology, Razi University,
Bagh-e Abrisham, Kermanshah, Iran

Ali Kosh on Deh Luren plain ($32^{\circ}33'28''\text{N}$, $47^{\circ}19'30''\text{E}$) was excavated in 1961 and 1963 by an American expedition directed by Frank Hole and Kent Flannery as a part of a larger regional project (Hole et al. 1969). Dated to c. 9500–8500 cal. BP (Hole 2000), Ali Kosh is an important Neolithic site located at the eastern edge of the Fertile Crescent (**Figure 1**), providing ample evidence of plant (cf. Moore 1982) and animal domestication (cf. Zeder 1999). In May 2017 a small stratigraphical trench was opened by Hojjat Darabi with the intention of revising the chronology of the site and to gather samples for research on subsistence strategies (Darabi 2017) (**Figure 2**).




Figure 1. Map of Deh Luren plain showing the location of Ali Kosh.

Drawing by S. Bahramiyan.

During this recent small-scale excavation, a dense cluster of 13 human burials was found at a depth of roughly 4–5m below the surface. Due to time constraints they were not explored *in situ*. Several of the skeletons were cut with large blocks of



Figure 2. A general view of the 2017 excavations at Ali Kosh.

surrounding soil and transported to the archaeological workshop at Razi University in Kermanshah. There they were cleaned and documented with a set of photographs enabling 3D modelling of each block. Finally, all human elements and artifacts were retrieved, cleaned and the human remains analyzed following the protocols of Buikstra and Ubelaker (1994).

In total, three blocks contained the remains of at least 11 individuals (**Figure 3**), although many of them were represented only by crania and/or mandibles while postcranial elements were under-represented. More specifically, block A contained one partially preserved skeleton (H2) and one cranium (H1), block B contained a cranium (H3) on top of three incomplete and disarticulated skeletons (H10, H11 and H12), another disarticulated cranium (H4), incomplete skeleton (H5) and disarticulated mandible (H7). The last block C included a skull (H6) with small remains of postcranial skeleton. Additionally, two mandibles (H8 and H9) were retrieved from other contexts and several dozen fragmented human elements have been identified in assemblages of animal remains. Additionally, one more skull excavated by Hole and Flannery in 1963 has been studied in the National Museum of Iran, referred to here as H13. Another skull curated in this museum (Niknami et al. 2011) was not available for study due to its use in an ongoing exhibition.



Figure 3. Human remains from Ali Kosh, soil block B during cleaning.

All blocks were affected by insect tunneling with a few nests observed, although bones were only slightly damaged by what may be attributed to solitary wasps (Pittoni 2009). In general, human remains were relatively well preserved in hard clayish soil with the only exception of inner parts of the crania that in most cases were empty and therefore strongly weathered and incrustated by crystalline deposits of various size (cf. Sołtysiak & Fetner 2017). All crania were distorted postmortem by pressure from the burial matrix resulting in element dislocations.

Although the burials were not excavated in a systematic way, several artifacts were retrieved from the soil surrounding the human elements. In particular, two subadult individuals received several hundred beads: around the pelvis of H5 there were shell beads, a bigger green one close to the face and two fine big mollusc shells close to the elbow and right auditory meatus. Around the pelvis of individual H2 small shells were found (Figure 4). Finally, a fragment of a fine narrow blade and a few small shell beads were located close to the bones of H6. Single small shells and fragments of flint tools were also scattered elsewhere. Most skeletons were coated with red ochre.

Only limited insight into the funerary customs is possible due to the accidental character of the discovery. The bodies of all three individuals with preserved articulations (H2, H5, H6) were buried in squatting position, both limbs hyperflexed, hands and knee joints close to face, feet and elbows close to pelvis. Such position perhaps involved some bundling with textiles or ropes, as all long bones were set together in vertical position between the skull and the os coxae. There were also many disarticulated bones and teeth, which were most likely the result of long-term use



Figure 4. Small mollusc shells around pelvic bones of individual H2.

of the cemetery, with some burials opened after complete skeletonization and bones re-buried in secondary contexts. Primary burials in squatting position seem to be not uncommon during the Pre-Pottery and Early Pottery Neolithic in the Near East (cf. Ortiz et al. 2013; Akkermans et al. 2006) and were also previously reported at Ali Kosh (Hole et al. 1969).

The available assemblage includes 7 crania, 7 mandibles and postcranial elements from at least 7 individuals (Table 1). The most striking feature of all crania was their more or less pronounced artificial deformation that was evident in spite of post-mortem alteration and fragmentation of all crania. In all cases circumferential modification was evident (Frieß & Baylac 2003), resulting from application of a band wrapped around the cranium along the anterior parietal and occipital, forming a conical protuberance around lambda (Figure 5). In most cases crania were very elongated and only in H3 the shape had been only slightly modified, still with clear impressions of the band. Change of cranium geometry affected the skull base, especially the occipital condyles, which were very convex, with the anterior part at a high angle to the horizontal plane. Additional crests between inferior nuchal line and foramen magnum were present in at least two individuals.

Artificial cranial deformation was common in the Near East and especially in Iran during the Neolithic and Chalcolithic (Meiklejohn et al. 1992; Daems & Croucher 2007), although its pattern changed over time. Among crania excavated by Hole and Flannery at least three female ones were artificially deformed (Hole et al. 1969), although in the female skull curated in the National Museum of Iran no clear evidence of deformation may be seen (Niknami et al. 2011). Artificial cranial deformation was



Figure 5. Cranial deformation in individual H6.

Table 1. Basic characteristics of skeletons from Ali Kosh. Cran. – cranium, Mand. – mandible, PCr – postcranial skeleton, URI1 Avul – avulsion of the right upper incisor, Calc – dental calculus. Scale for completeness: 0 – absent, 1 – less than 50% preserved, 2 – more than 50% preserved, 3 – complete or nearly complete.

ID	Sex	Age-at-death	Completeness			Caries	URI1 Avul	Enamel hypoplasia			Calc
			Cran.	Mand.	PCr			C	M1	Other	
H1	F??	old	3	0	0	0/1	?				+
H2		9	3	1	2	0/5	-	+	+	+	-
H3	F??	15	3	0	0	0/3	-	-	-	-	-
H4	M??	young	3	0	0	0/7	+	-	-	-	+
H5		6	3	3	2	0/11	-	-	-	-	-
H6	M	young	3	3	1	0/28	+	+	-	-	+
H7	M??	young	0	2	0	0/5			-	-	-
H8	M??	young	0	2	0	0/9			-	-	-
H9		7	0	2	1	0/4		+	+	+	-
H10	M??	adult	0	0	1						
H11	F??	14-21	0	0	1						
H12		7-14	0	0	1						
H13	M?	adult	2	0	0	0/6	+		-	+	-

also observed at the nearby site of Chogha Sefid (Hole 1977). The presence of this custom in the recently excavated assemblage at Ali Kosh is, as such, not surprising.

Another cultural modification of the head observed at Ali Kosh was avulsion of the upper right first incisor in all adult males, but not in children nor adolescent individuals. Small sample size makes it impossible to say whether females were also affected, as the only adult female cranium in the sample belonged to an old individual with many teeth (incisors included) lost antemortem. However, in three male individuals even if several teeth were lost postmortem, evidence of antemortem avulsion of this one specific tooth was evident. As two children and one adolescent were not afflicted, it is possible that this custom of tooth avulsion served as an age related rite of passage. Tooth avulsion was common during the Early Holocene in North Africa (Stojanowski et al. 2014; De Groote & Humphrey 2016), and it was also occasionally observed in the Natufian culture, with no relation to sex or age-at-death (Boyd 2002). Such a custom has not been previously reported in the eastern part of the Fertile Crescent.

The age-at-death profile for the Ali Kosh sample seems to be highly biased; there are no infants and only one old adult and therefore age categories with the highest mortality rate are underrepresented, even when small sample size is taken into account. No cases of dental caries have been observed, suggesting a possible diet with low intake of fermentable sugars (cf. Soltysiak 2014). Frequency of cranial porosities was also rather low, with only one case (H6) of highly obliterated microporosity at several locations on the cranium. On the other hand, in two individuals (H1 and H3, both probable females) the temporal line was very pronounced, with clear vascularization above, suggesting some mechanical stress on the temporal muscle.

Linear enamel hypoplasia (LEH) follows an unusual pattern in this assemblage. Many individuals were free or almost free from this condition, but in two subadult dentitions (H2 and H9) multiple LEH were present in the canines, also along the upper part of the crown. The regular pattern of LEH lines suggests a seasonal character of the stress episodes. Along with the canines, a single hypoplastic line was observed also on the first molars, close to their cusps. Such a condition is very unusual and suggests an episode of stress in infancy around the first year of life. Also in one adult individual (H6) seasonally patterned LEH was observed on the canines. Although interpretation of LEH is difficult because of the osteological paradox (Wood et al. 1992), the pattern of LEH at Ali Kosh is very specific and different from what is seen at later sites where this condition is present in higher frequency, but hypoplastic lines are usually observed mainly close to the cemento-enamel junction on the anterior teeth (cf. Soltysiak 2010).

Acknowledgements. Research on human remains from Ali Kosh was financially supported by the Polish National Science Centre (grant No. 2016/22/M/HS3/00353)

and by Razi University in Kermanshah. Many thanks to Dr. Feridoun Biglari for making the skull stored in the National Museum of Iran available for the present research, and to Hafez Ghaderi and Saman Mostafapour for their valuable help in cleaning bones from very hard soil.

References

- Akkermans P.M.M.G., Cappers R., Cavallo C., Nieuwenhuys O., Nilhamn B., Otte I.N. (2006), *Investigating the Early Pottery Neolithic of Northern Syria: New evidence from Tell Sabi Abyad*, *American Journal of Archaeology* 110(1):123–156.
- Boyd B. (2002), *Ways of eating/ways of being in the Later Epipalaeolithic (Natufian) Levant* [in:] “Thinking through the body,” Y. Hamilakis, M. Pluciennik, S. Tarlow (eds.), Boston: Springer, pp. 137–152.
- Buikstra J.A., Ubelaker D.H. (Eds.) (1994), *Standards for data collection from human skeletal remains*, Fayetteville: Arkansas Archaeological Survey.
- Daems A., Croucher K. (2007), *Artificial cranial modification in prehistoric Iran: Evidence from crania and figurines*, *Iranica Antiqua* 42:1–21.
- Darabi H. (2017), *A report on the stratigraphy and chronology of Tapeh Ali Kosh*, Tehran: Iranian Center for Archaeological Research. (in Farsi)
- De Groote I., Humphrey L.T. (2016), *Characterizing evulsion in the Later Stone Age Maghreb: Age, sex and effects on mastication*, *Quaternary International* 413:50–61.
- Frieß M., Baylac M. (2003), *Exploring artificial cranial deformation using elliptic Fourier analysis of procrustes aligned outlines*, *American Journal of Physical Anthropology* 122(1):11–22.
- Hole F. (1977), *Studies in the archeological history of the Deh Luran Plain: The excavation of Chagha Sefid*, Ann Arbor: University of Michigan.
- Hole F. (2000), *New radiocarbon dates for Ali Kosh, Iran*, *Neo-Lithics* 1:13.
- Hole F., Flannery K.V., Neely J.A. (1969), *Prehistory and human ecology of the Deh Luran plain. An early village sequence from Khuzistan, Iran*, Ann Arbor: University of Michigan.
- Meiklejohn C., Agelarakis A., Akkermans P.M.M.G., Smith P.E.L., Solecki R. (1992), *Artificial cranial deformation in the Proto-Neolithic and Neolithic Near East and its possible origin: Evidence from four sites*, *Paléorient* 18(2):83–97.
- Moore A.M.T. (1982), *Agricultural origins in the Near East: A model for the 1980s*, *World Archaeology* 14(2):224–236.
- Niknami K.A., Ramazani M., Niknami N. (2011), *Ali Kosh Lady and her artificially modified head: An appraisal*, *Iranian Journal of Archaeological Studies* 1(2):17–24.
- Ortiz A., Chambon P., Molist M. (2013), “Funerary bundles” in the PPNB at the archaeological site of Tell Halula (middle Euphrates valley, Syria): Analysis of the

- taphonomic dynamics of seated bodies*, *Journal of Archaeological Science* 40(12): 4150–4161.
- Pittoni E. (2009), *Necropoli of Pill'e Matta Quartucciu (Cagliari, Sardinia): Wild bee and solitary wasp activity and bone diagenetic factors*, *International Journal of Osteoarchaeology* 19(3):386–396.
- Sołtysiak A. (2010), *Death and decay at the dawn of the city. Interpretation of human bone deposits at Tell Majnuna, Areas MTW, EM and EMS*, Warszawa: Instytut Archeologii UW.
- Sołtysiak A. (2014), *Frequency of dental caries as a proxy indicator of mobility: The case of the Khabur basin human populations* [in:] “Paleonutrition and food practices in the ancient Near East. Towards a multidisciplinary approach,” L. Milano (ed.), Padova: S.A.R.G.O.N. Editrice e Libreria, pp. 53–70.
- Sołtysiak A., Fetner R.A. (2017), *Taphonomy of human remains exposed in burial chambers, with special reference to Near Eastern hypogea, ossuaries and burial caves* [in:] “Engaging with the dead: Exploring changing human beliefs about death, mortality and the human body,” J. Bradbury, C. Scarre (eds.), Oxford & Havertown: Oxbow Books, pp. 67–76.
- Stojanowski C.M., Carver C.L., Miller K.A. (2014), *Incisor avulsion, social identity and Saharan population history: New data from the Early Holocene southern Sahara*, *Journal of Anthropological Archaeology* 35:79–91.
- Wood J.W., Milner G.R., Harpending H.C., Weiss K.M. (1992), *The osteological paradox: Problems of inferring prehistoric health from skeletal samples*, *Current Anthropology* 33(4):343–370.
- Zeder M. (1999), *Animal domestication in the Zagros: A review of past and current research*, *Paléorient* 25(2):11–25.